

2015

Heart Failure and Depression: Reducing Readmission Rates

Janelle Maciej

CentraCare Health, maciejj@centracare.com

Follow this and additional works at: https://digitalcommons.centracare.com/nursing_posters



Part of the [Other Nursing Commons](#)

Recommended Citation

Maciej, Janelle, "Heart Failure and Depression: Reducing Readmission Rates" (2015). *Nursing Posters*. 42.
https://digitalcommons.centracare.com/nursing_posters/42

This Book is brought to you for free and open access by the Posters and Scholarly Works at DigitalCommons@CentraCare Health. It has been accepted for inclusion in Nursing Posters by an authorized administrator of DigitalCommons@CentraCare Health. For more information, please contact schlepers@centracare.com.



Heart Failure and Depression: Reducing Readmission Rates



Janelle Maciej, RN ,BSN, BC

Institution, St. Cloud, Minnesota

Purpose Statement

Reduce heart failure readmissions through implementation of a depression assessment in the acute care setting.

Synthesis of Evidence

Reviewed over 40 articles

- Patients with major depression were readmitted twice as often as non-depressed patients. Grade A.
- Major depression in hospitalized patients was associated with an increased mortality at 3 months and at 1 year. Grade B.
- There is a strong association between depression and the severity of heart failure. Grade B.
- SCH readmission rate cost is about \$10,697 per readmitted patient.
- PHQ-2 depression screen has high specificity (0.91), has high negative predictive value(0.87) and low sensitivity(0.52). Grade B.

Team Members

- Angie Lieser RN
Telemetry Charge Nurse
- Mary Schiminich RN
Heart Failure Specialist
- Ryan Zinken RN
Behavioral Health Case Manager
- Daren Hendrickson RN
Behavioral Health Case Manager

EBP Practice Change

- MD will order Heart Failure order set.

Other Nursing Orders (Through next 24h)

Start

11/04/14 1646 [Nursing to administer the PHQ-2 depression screening tool.](#) [139410101] FYI [Complete](#) [Discontinue](#)

Order Set: Adult Admission Orders (23)

- The Patient Health Questionnaire-2(PHQ-2) will automatically populate on the Physical Assessment flow

PHQ-2 Depression Screening. Over the last 2 weeks, how often have you been bothered by any of the following problems? *Complete once during admission*

Little interest or pleasure in doing	2			
Feeling down, depressed or	2			
Total PHQ-2 Score	4			

- Depression screen will populate on the Work List

Complete PHQ 2 Routine Doc Skip

- If positive, a Best Practice Advisory (BPA) will fire for a Behavioral Health Case Manager (BHCM) consult.

BestPractice Advisory - Tele,Adam

Positive PHQ-2 depression screen - Please place a Behavioral Health Case Manager Consult and include reason for consult.

Accept Cancel

- The BHCM will interview the patient and then recommend the best next steps. Follow up with Primary MD, a Psychiatrist consult, Psychology consult etc.



Follow Up Appointments

Holmgren, Leah A, MD
Specialty: Internal Medicine

Appointments/Instructions: Heart Failure hospitalization and positive depression screen

- There may be "Depression Screening" in the epic problem list. **Hospital Problem List**
Depression screen
- ICD-9-CM:
V79.0
ICD-10-CM:
Z13.89

Pre/Post Measures



References

Elderson, Smolderen, et al (2011). Accuracy and Prognostic value of American Heart Association-Recommended Depression Screening in Patients with Coronary Heart Disease: Data From The Heart and Soul Study. *Journal of the American Heart Association* 4, 533-540.

Freeland, Rich et al (2003). Prevalence of Depression in Hospitalized Patients with Congestive Heart Failure *Psychosomatic Medicine* 65, 119-128.

Jiang, Kuchibhatla et al (2007). Relationship between depressive symptoms and long-term mortality in patients with heart failure. *American Heart Journal* 154(1), 102-108.

Jiang, Alexander et al (2001). Relationship of Depression to Increased risk of Mortality and Rehospitalization in Patients with Congestive Heart Failure. *Arch Intern Med*, 161, 1849-1856.

Rollman, Belnap et al (2012). A positive phq-2 depression screen among hospitalized heart failure patients is associated with elevated 12- month mortality *J card fail* 18(3), 238-245.

Rutledge, Reis et al (2006). Depression in Hear Failure a Meta analytic review of prevalence, intervention effects and associated with clinical outcomes. *Journal of the American College of Cardiology* 48(8),1527-1537.

Sowden, Mastroiuro et al (2010). Detection of depression in cardiac inpatients: Feasibility and results of systematic screening *American Heart Journal* 159(5), 780-787.